



Ferric Phosphate (hydrated) 2,97 % w/w  
(equivalent to anhydrous FePO4: min 21,1 g/kg)  
Granular ready-made bait (RB)

# SLUXX HP

Granular snailicide bait for the control of snails and slugs. It has an attractive action and acts on the stomach. After ingestion of the product, snails and slugs stop feeding and secrete secretions into plants. There is no need to collect the dead snails because after receiving the bait they tend to go to isolated places to die.

SCOPE	TARGET	DOSES	Way and time application	Maximum number of applications per growing season
		Kg/acre (max)		
All crops edible and non-edible (G + F)	Snails and Slugs	0,7	Uniform scattering in the field. Immediately after sowing or planting or with the appearance of the infestation or with the first symptoms in the plants.	4



**Remarks:** 1. Prefer preventive applications or apply as soon as the first damages appear. 2. The product can be used at all stages of plant development. 3. Repeat application every 2 weeks, if necessary or after heavy rains or when the amount of preparation in the field has been significantly reduced. Repeat application when the seeds of the preparation have been consumed and the weather conditions are favorable for snails or slugs until the end of cultivation. 4. Only tou is used. 5. Do not apply it against the wind. 6. Its action is not affected by rain or wet conditions.

**Method of application:** Spread evenly over the entire surface of the field or crop, taking care not to leave grains of the preparation in the foliage of the plants. Apply with a granular fertilizer dispenser, preferably in the early afternoon, on moist soil after rain or watering.

**Special agricultural, phytosanitary or environmental conditions under which the preparation can be used or excluded:** Contact of grains with sensitive leaves (eg sprouts in spring crops, greenhouses or under cover) may cause burns or local necrosis.

**Evidence of phytotoxicity, variety susceptibility and any other side effects on plants or their products:** No phytotoxicity at recommended uses and doses.

**Safety period between application and (i) sowing or planting of the protected crop: -, (ii) sowing or planting of the following crops: -, (iii) human or animal access to the crop to which the preparation has been applied: --.**

**Last intervention before harvest or before placing on the market in the case of post-harvest uses: all crops: --.**

**Storage conditions, time stability of the preparation:** Store sealed in its original packaging, in a dry, cool place, without the effect of direct sunlight and extreme temperatures. In these conditions it remains stable for two (2) years from the date of manufacture.

**PACKAGING:** 350gr / 500gr / 700gr / 1Kg / 1,4Kg / 5Kg / 7Kg / 15Kg / 20Kg / 25Kg

**THE INFORMATION WRITTEN IN THIS FORM HAS AN INFORMATIONAL CHARACTER AND DOES NOT SUBSTITUTE IN ANY CIRCUMSTANCES. PLANT PROTECTION PRODUCTS ARE INTENDED FOR USE BY PROFESSIONAL USERS. INSTRUCTIONS FOR USE AND PRECAUTIONS, WRITTEN ON THE LABEL MUST BE OBSERVED. FOR MORE INFORMATION CONTACT THE TECHNICAL DEPARTMENT OF OUR COMPANY OR CONSULT THE LOCAL AGRONOMISTS.**

NitroFarm SA is certified with EN ISO 9001:2015 Quality Management System and EN ISO 14001:2015 Environmental Management System

**bio**

suitable for use in organic farming, according to the requirements of EU Regulation 848/2018 & 1165/2021

**NITROFARM**

134 07/2024